



COL9A2 gene

collagen type IX alpha 2 chain

Normal Function

The *COL9A2* gene provides instructions for making part of a large molecule called type IX collagen. Collagens are a family of proteins that strengthen and support connective tissues, such as skin, bone, cartilage, tendons, and ligaments. In particular, type IX collagen is an important component of cartilage, which is a tough, flexible tissue that makes up much of the skeleton during early development. Most cartilage is later converted to bone, except for the cartilage that continues to cover and protect the ends of bones and is present in the nose and external ears.

Type IX collagen is made up of three proteins that are produced from three distinct genes: one $\alpha 1$ (IX) chain, which is produced from the *COL9A1* gene, one $\alpha 2$ (IX) chain, which is produced from the *COL9A2* gene, and one $\alpha 3$ (IX) chain, which is produced from the *COL9A3* gene. Type IX collagen is more flexible than other types of collagen molecules and is closely associated with type II collagen. Researchers believe that the flexible nature of type IX collagen allows it to act as a bridge that connects type II collagen with other cartilage components. Studies have shown that type IX collagen also interacts with the proteins produced from the *MATN3* and *COMP* genes.

Health Conditions Related to Genetic Changes

intervertebral disc disease

multiple epiphyseal dysplasia


At least five mutations in the *COL9A2* gene have been shown to cause dominant multiple epiphyseal dysplasia, a disorder of cartilage and bone development that primarily affects the ends of the long bones in the arms and legs (epiphyses). All of these mutations disrupt how genetic information is spliced together to make the blueprint for producing the $\alpha 2$ (IX) chain. These mutations, called splice-site mutations, change one DNA building block (nucleotide) near an area of the gene called exon 3. These mutations in the *COL9A2* gene result in the deletion of 12 protein building blocks (amino acids) from the $\alpha 2$ (IX) chain. It is not known how mutations in *COL9A2* cause the signs and symptoms of dominant multiple epiphyseal dysplasia.

Stickler syndrome

Chromosomal Location

Cytogenetic Location: 1p34.2, which is the short (p) arm of chromosome 1 at position 34.2

Molecular Location: base pairs 40,300,487 to 40,317,653 on chromosome 1 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- alpha 2 type IX collagen
- CO9A2_HUMAN
- collagen IX, alpha-2 polypeptide
- collagen type IX alpha 2
- collagen, type IX, alpha 2
- EDM2
- epiphyseal dysplasia, multiple 2

Additional Information & Resources

Educational Resources

- Molecular Biology of the Cell (fourth edition, 2002): Collagens Are the Major Proteins of the Extracellular Matrix
<https://www.ncbi.nlm.nih.gov/books/NBK26810/#A3551>
- Molecular Cell Biology (fourth edition, 2000): Collagen: The Fibrous Proteins of the Matrix
<https://www.ncbi.nlm.nih.gov/books/NBK21582/>

GeneReviews

- Multiple Epiphyseal Dysplasia, Dominant
<https://www.ncbi.nlm.nih.gov/books/NBK1123>
- Stickler Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1302>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28COL9A2%5BTIAB%5D%29+OR+%28EDM2%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+2160+days%22%5Bdp%5D>

OMIM

- COLLAGEN, TYPE IX, ALPHA-2
<http://omim.org/entry/120260>
- INTERVERTEBRAL DISC DISEASE
<http://omim.org/entry/603932>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_COL9A2.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=COL9A2%5Bgene%5D>
- HGNC Gene Family: Collagen proteoglycans
<http://www.genenames.org/cgi-bin/genefamilies/set/575>
- HGNC Gene Family: Collagens
<http://www.genenames.org/cgi-bin/genefamilies/set/490>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=2218
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/1298>
- UniProt
<http://www.uniprot.org/uniprot/Q14055>

Sources for This Summary

- Baker S, Booth C, Fillman C, Shapiro M, Blair MP, Hyland JC, Ala-Kokko L. A loss of function mutation in the COL9A2 gene causes autosomal recessive Stickler syndrome. *Am J Med Genet A*. 2011 Jul;155A(7):1668-72. doi: 10.1002/ajmg.a.34071. Epub 2011 Jun 10.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21671392>
- Briggs MD, Chapman KL. Pseudoachondroplasia and multiple epiphyseal dysplasia: mutation review, molecular interactions, and genotype to phenotype correlations. *Hum Mutat*. 2002 May; 19(5):465-78. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11968079>
- GeneReview: Multiple Epiphyseal Dysplasia, Dominant
<https://www.ncbi.nlm.nih.gov/books/NBK1123>
- Higashino K, Matsui Y, Yagi S, Takata Y, Goto T, Sakai T, Katoh S, Yasui N. The alpha2 type IX collagen tryptophan polymorphism is associated with the severity of disc degeneration in younger patients with herniated nucleus pulposus of the lumbar spine. *Int Orthop*. 2007 Feb;31(1):107-11. Epub 2006 Apr 4.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16586133>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2267527/>
- Jim JJ, Noponen-Hietala N, Cheung KM, Ott J, Karppinen J, Sahraravand A, Luk KD, Yip SP, Sham PC, Song YQ, Leong JC, Cheah KS, Ala-Kokko L, Chan D. The TRP2 allele of COL9A2 is an age-dependent risk factor for the development and severity of intervertebral disc degeneration. *Spine (Phila Pa 1976)*. 2005 Dec 15;30(24):2735-42.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16371896>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/COL9A2>

Reviewed: January 2013
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services